



Gulf of Mexico Harmful Algal Bloom Bulletin

Region: Southwest Florida


Monday, 28 April 2014

NOAA National Ocean Service

NOAA Satellite and Information Service

NOAA National Weather Service

Last bulletin: Monday, April 21, 2014



Satellite chlorophyll image with possible *K. brevis* HAB areas shown by red polygon(s), when applicable. Points represent cell concentration sampling data from April 18 to 24: red (high), orange (medium), yellow (low b), brown (low a), blue (very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Florida Fish and Wildlife Conservation Commission (FWC) Fish and Wildlife Research Institute. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

http://tidesandcurrents.noaa.gov/hab/habfs_bulletin_guide.pdf

Detailed sample information can be obtained through FWC Fish and Wildlife Research Institute at:

<http://myfwc.com/redtidestatus>

To see previous bulletins and forecasts for other Harmful Algal Bloom Bulletin regions, visit at: <http://tidesandcurrents.noaa.gov/hab/bulletins.html>

Conditions Report

There is currently no indication of *Karenia brevis* (commonly known as Florida red tide) along the coast of southwest Florida, including the Florida Keys. No respiratory irritation is expected Monday, April 28 through Monday, May 5.

Check http://tidesandcurrents.noaa.gov/hab/beach_conditions.html for recent, local observations.


Analysis

Samples collected over the last week along the coast of southwest Florida from Pinellas to Collier counties all indicate that *Karenia brevis* is not present, with the exception of two samples that identified background concentrations in the bay regions of northern Sarasota at New Pass and Bay Dock (FWRI, MML, CCPCPD, SCHD; 4/18-4/23).

In recent MODIS Aqua imagery (4/25, shown left) patches of elevated chlorophyll (2-10 $\mu\text{g/L}$) are visible along- and offshore portions of the southwest Florida coast from Pinellas to Collier counties.

Harmful algal bloom formation at the coast of southwest Florida is not expected today through Monday, May 5.


Burrows, Kavanaugh



Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS).

Wind Analysis

Southwest Florida: Southeast winds (10-15kn, 5-8m/s) becoming southwest in the afternoon. South to southeast winds (5-15kn, 3-8m/s) tonight through Thursday becoming southwest to west winds (5-10kn, 3-5m/s) Thursday afternoon through evening. Northeast winds (10kn, 5m/s) after midnight becoming northwest to north winds (5-15kn) Friday.



Satellite chlorophyll image and forecast winds for April 29, 2014 06Z with points representing cell concentration sampling data from April 18 to 24: red (high), orange (medium), yellow (low b), brown (low a), blue (very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Florida Fish and Wildlife Conservation Commission (FWC) Fish and Wildlife Research Institute. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

http://tidesandcurrents.noaa.gov/hab/habfs_bulletin_guide.pdf

Verified and suspected HAB areas shown in red. Other areas of high chlorophyll concentration shown in yellow (see p. 1 analysis for interpretation).