



Gulf of Mexico Harmful Algal Bloom Bulletin

Region: Southwest Florida


Tuesday, 20 January 2015

NOAA National Ocean Service

NOAA Satellite and Information Service

NOAA National Weather Service

Last bulletin: Monday, January 12, 2015



Satellite chlorophyll image with possible *K. brevis* HAB areas shown by red polygon(s), when applicable. Points represent cell concentration sampling data from January 11 to 19: red (high), orange (medium), yellow (low b), brown (low a), blue (very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Florida Fish and Wildlife Conservation Commission (FWC) Fish and Wildlife Research Institute. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

http://tidesandcurrents.noaa.gov/hab/habfs_bulletin_guide.pdf

Detailed sample information can be obtained through FWC Fish and Wildlife Research Institute at:

<http://myfwc.com/redtidestatus>

To see previous bulletins and forecasts for other Harmful Algal Bloom Bulletin regions, visit at: <http://tidesandcurrents.noaa.gov/hab/bulletins.html>

Conditions Report

There is currently no indication of *Karenia brevis* (commonly known as Florida red tide) along the coast of southwest Florida, including the Florida Keys. No respiratory irritation is expected alongshore southwest Florida, including the Florida Keys, Tuesday, January 20 through Monday, January 26.

Check http://tidesandcurrents.noaa.gov/hab/beach_conditions.html for recent, local observations.


Analysis

The most recent samples received from alongshore southwest Florida, from Pinellas to Lee counties, and along- and offshore the Gulf side and ocean side of the Lower and Middle Florida Keys, all indicate that *Karenia brevis* is not present (FWRI, MML, SCHD; 1/12-14).

Recent MODIS Aqua imagery (1/19, shown left) is partially obscured by clouds along- and offshore southwest Florida from Sarasota County to the Florida Keys, limiting analysis. In southwest Florida, elevated chlorophyll (2-8 $\mu\text{g/L}$) is visible along the coast from Pinellas County to southern Lee County.

Harmful algal bloom formation at the coast of southwest Florida is not expected today through Monday, January 26.


Davis, Kavanaugh



Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS).

Wind Analysis

Englewood to Tarpon Springs (Venice): South winds (5-10kn, 3-5m/s) today becoming east winds after midnight. East winds (5kn, 3m/s) Wednesday becoming north winds (5-10kn) in the afternoon and then becoming northeast after midnight. East to northeast winds (5-10kn) Thursday. Southeast to south winds (10kn, 5m/s) Friday becoming southwest (10kn) Friday evening. West to northwest winds (10-15kn, 5-8m/s) Saturday.



Satellite chlorophyll image and forecast winds for January 21, 2015 06Z with points representing cell concentration sampling data from January 11 to 19: red (high), orange (medium), yellow (low b), brown (low a), blue (very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Florida Fish and Wildlife Conservation Commission (FWC) Fish and Wildlife Research Institute. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

http://tidesandcurrents.noaa.gov/hab/habfs_bulletin_guide.pdf

Verified and suspected HAB areas shown in red. Other areas of high chlorophyll concentration shown in yellow (see p. 1 analysis for interpretation).